

AMENDMENTS TO THE CLAIMS

1. (Currently amended) A method for treating inflammatory disease of the external segment or the anterior segment of the eye, which comprises administering to the eye an aqueous eye drop comprising 2-amino-3-(4-bromobenzoyl)phenylacetic acid or its pharmacologically acceptable salt or a hydrate thereof and at least one organic amine selected from the group consisting of an amino acid, an alkanolamine, ~~ethylenediamine, trimethylenediamine,~~ a piperazine and an aminoalkylsulfonic acid once a day, and maintaining a therapeutically effective concentration of 2-amino-3-(4-bromobenzoyl)phenylacetic acid in the anterior aqueous humor of the eye ~~for at least 24 hours after the intraocular administration.~~

2. (Cancelled)

3. (Previously presented) The method according to claim 1, wherein the concentration of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or its pharmacologically acceptable salt or a hydrate thereof in the aqueous eye drop is 0.01 to 0.5w/v%.

4. (Cancelled)

5. (Previously presented) The method according to claim 1, wherein the organic amine is an amino acid and its concentration is 0.35 to 5w/v%.

6. (Previously presented) The method according to claim 1, wherein the organic amine is an alkanolamine and its concentration is 0.15 to 0.95w/v%.

7. (Previously presented) The method according to claim 1, wherein the alkanolamine is trometamol.

8. (Cancelled)

9. (Previously presented) The method according to claim 1, wherein the organic amine is a piperazine and it is contained at a concentration of 0.05 to 5w/v% in the aqueous eye drop.

10. (Previously presented) The method according to claim 1, wherein the organic amine is an aminoalkylsulfonic acid and its concentration is 0.05 to 5w/v%.

11. (Previously presented) The method according to claim 10, wherein the aminoalkylsulfonic acid is aminoethylsulfonic acid.

12-35. (Cancelled)